

Publikationen zu TDM von Antibiotika im Labor Dr. Brunner

Ehmann L, Zoller M, Minichmayr IK, Scharf C, Huisinga W, **Zander J**, Kloft C. Development of a dosing algorithm for meropenem in critically ill patients based on a population pharmacokinetic/pharmacodynamic analysis. *Int J Antimicrob Agents*. 2019;54:309-317.

Li X, Zoller M, Fuhr U, Huseyn-Zada M, Maier B, Vogeser M, **Zander J**, Taubert M. Ciprofloxacin in critically ill subjects: considering hepatic function, age and sex to choose the optimal dose. *J Antimicrob Chemother*. 2019;74:682-690.

Ehmann L, Zoller M, Minichmayr IK, Scharf C, Maier B, Schmitt MV, Hartung N, Huisinga W, Vogeser M, Frey L, **Zander J**, Kloft C. Role of renal function in risk assessment of target non-attainment after standard dosing of meropenem in critically ill patients: a prospective observational study. *Crit Care*. 2017;21:263.

Taubert M, **Zander J**, Frechen S, Scharf C, Frey L, Vogeser M, Fuhr U, Zoller M. Optimization of linezolid therapy in the critically ill: the effect of adjusted infusion regimens. *J Antimicrob Chemother*. 2017;72:2304-2310.

Lucha S, Taibon J, Pongratz S, Geletneky C, Huber E, Wintterle-Roehm C, Lang R, Grimm SH, Duelffer T, Tarasov K, **Zander J**, Vogeser M, Kobold U. An LC-MS/MS based candidate reference method for the quantification of total gentamicin in human serum and plasma using NMR characterized calibrator material. *Clin Chim Acta*. 2017;464:211-217.

Zander J, Döbbeler G, Nagel D, Scharf C, Huseyn-Zada M, Jung J, Frey L, Vogeser M, Zoller M. Variability of piperacillin concentrations in relation to tazobactam concentrations in critically ill patients. *Int J Antimicrob Agents*. 2016;48:435-9. doi:

Taubert M, Zoller M, Maier B, Frechen S, Scharf C, Holdt LM, Frey L, Vogeser M, Fuhr U, **Zander J**. Predictors of Inadequate Linezolid Concentrations after Standard Dosing in Critically Ill Patients. *Antimicrob Agents Chemother*. 2016;60:5254-61.

Zander J, Döbbeler G, Nagel D, Maier B, Scharf C, Huseyn-Zada M, Jung J, Frey L, Vogeser M, Zoller M. Piperacillin concentration in relation to therapeutic range in critically ill patients--a prospective observational study. *Crit Care*. 2016;20:79.

Zander J, Maier B, Zoller M, Döbbeler G, Frey L, Teupser D, Vogeser M. Effects of biobanking conditions on six antibiotic substances in human serum assessed by a novel evaluation protocol. *Clin Chem Lab Med*. 2016;54:265-74.

Zoller M, Döbbeler G, Maier B, Vogeser M, Frey L, **Zander J**. Can cytokine adsorber treatment affect antibiotic concentrations? A case report. *J Antimicrob Chemother*. 2015;70:2169-71.

Zander J, Maier B, Suhr A, Zoller M, Frey L, Teupser D, Vogeser M. Quantification of piperacillin, tazobactam, cefepime, meropenem, ciprofloxacin and linezolid in serum using an isotope dilution UHPLC-MS/MS method with semi-automated sample preparation. *Clin Chem Lab Med*. 2015;53:781-91.

Zoller M, Maier B, Hornuss C, Neugebauer C, Döbbeler G, Nagel D, Holdt LM, Bruegel M, Weig T, Grabein B, Frey L, Teupser D, Vogeser M, **Zander J**. Variability of linezolid concentrations after standard dosing in critically ill patients: a prospective observational study. *Crit Care*. 2014;18:R148.

Zander J, Bruegel M, Kleinhempel A, Becker S, Petros S, Kortz L, Dorow J, Kratzsch J, Baber R, Ceglarek U, Thiery J, Teupser D. Effect of biobanking conditions on short-term stability of biomarkers in human serum and plasma. *Clin Chem Lab Med*. 2014;52:629-39.

Zander J, Maier B, Zoller M, Teupser D, Vogeser M. Quantification of linezolid in serum by LC-MS/MS using semi-automated sample preparation and isotope dilution internal standardization. *Clin Chem Lab Med*. 2014;52:381-9.

König K, Kobold U, Fink G, Leinenbach A, Dülffer T, Thiele R, **Zander J**, Vogeser M. Quantification of vancomycin in human serum by LC-MS/MS. *Clin Chem Lab Med*. 2013;51:1761-9.